

# Asymptotic analysis of hedging errors in models with jumps

Peter Tankov

LPMA, Université Paris-Diderot — Paris 7

Based on joint work with E. Voltchkova (Université Toulouse 1)

Vienna, November 16, 2007

# Hedging in incomplete markets

- Incomplete market: exact replication impossible.
- Hedging is now an approximation problem.
- Industry practice: sensitivities to risk factors

Delta =  $\frac{\partial C(t, S_t)}{\partial S}$  : infinitesimal moves, hedge with stock

Gamma =  $\frac{\partial^2 C(t, S_t)}{\partial S^2}$  : bigger moves; hedge with liquid options

- Quadratic hedging: control the residual error

$$\min_{\phi} E \left( c + \int_0^T \phi_t dS_t - Y \right)^2$$

All these strategies require a continuously rebalanced portfolio.

# Discrete hedging

- Continuous rebalancing is unfeasible: in practice, the strategy  $\phi_t$  is replaced with a discrete strategy, leading to the hedging error of the “second type”: error of approximating the continuous portfolio with a discrete one.
- The simplest choice is  $\phi_t^n := \phi_{h[t/h]}$ ,  $h = T/n$ .
- This discretization error has only been studied in the case of continuous processes.

# Discrete hedging: the complete market case

- Bertsimas, Kogan and Lo '98 introduced an *asymptotic approach* allowing to study discrete hedging in continuous time.

Suppose

$$\frac{dS_t}{S_t} = \mu(t, S_t)dt + \sigma(t, S_t)dW_t$$

and we want to hedge a European option with payoff  $h(S_T)$  using delta-hedging  $\phi_t = \frac{\partial C}{\partial S}$ .

# CLT for hedging error

The discrete hedging error is defined by

$$\varepsilon_T^n = h(S_T) - \int_0^T \phi_t^n dS_t$$

Then  $\varepsilon_T^n \rightarrow 0$  but the renormalized error  $\sqrt{n}\varepsilon_T^n$  converges to

$$\sqrt{\frac{T}{2}} \int_0^T \frac{\partial^2 C}{\partial S^2} S_t^2 \sigma_t^2 dW_t^*,$$

where  $W^*$  is a Brownian motion independent of  $W$ .

- Hedging error decays as  $\sqrt{h}$ .
- It is orthogonal to the stock price.
- The amplitude is determined by the gamma  $\frac{\partial^2 C}{\partial S^2}$

# Intuition

In complete market,

$$\varepsilon_T^n = \int_0^T (\phi_t - \phi_t^n) dS_t$$

Let  $S_t = W_t$  and consider the renormalized error over one hedging interval:

$$\frac{1}{\sqrt{h}} \int_0^h \left( \frac{\partial C}{\partial S}(W_t) - \frac{\partial C}{\partial S}(0) \right) dW_t \approx \frac{1}{\sqrt{2}} \frac{\partial^2 C}{\partial S^2} \frac{1}{\sqrt{2h}} (W_h^2 - h)$$

The random variable  $\frac{1}{\sqrt{2h}} (W_h^2 - h)$  has mean zero, variance  $h$  and is uncorrelated with  $W_h$ .

# Approximating hedging portfolios

Hayashi and Mykland '05 interpreted the discrete hedging error as the error of approximating the “ideal” hedging portfolio  $\int_0^T \phi_t dS_t$  with a feasible hedging portfolio  $\int_0^T \phi_t^n dS_t$

- This makes sense in incomplete markets

Suppose  $\phi$  and  $S$  are Itô process:

$d\phi_t = \tilde{\mu}_t dt + \tilde{\sigma}_t dW_t$  and  $dS_t = \mu_t dt + \sigma_t dW_t$ . Then

$$\sqrt{n}\varepsilon_t^n \Rightarrow \sqrt{\frac{T}{2}} \int_0^t \tilde{\sigma}_s \sigma_s dW_s^*,$$

$$\text{where } \varepsilon_t^n := \int_0^t (\phi_t - \phi_t^n) dS_t.$$

- Weak convergence of processes in the Skorokhod topology on the space  $\mathbb{D}$  of càdlàg functions

# Discrete hedging in presence of jumps

The idea of approximating stochastic integrals goes back to Rootzen (80)

More recently, results by Geiss (02), (06), (07) but all authors work with continuous processes

Our idea: study the discretization error

$$\varepsilon_t^n := \int_0^t (\phi_{t-} - \phi_{t-}^n) dS_t$$

in presence of jumps in the underlying and the hedging strategy.

- Some tools are available in the study of the approximation error of the Lévy-driven Euler scheme by Jacod and Protter (98)

## Model setup: Lévy-Itô processes

$$X_t = X_0 + \int_0^t \mu_s ds + \int_0^t \sigma_s dW_s + \int_0^t \int_{|z| \leq 1} \gamma_s(z) \tilde{J}(ds \times dz) + \int_0^t \int_{|z| > 1} \gamma_s(z) J(ds \times dz).$$

- $J$ : Poisson random measure with intensity  $dt \times \nu$
- $\mu$  and  $\sigma$  are càdlàg  $(\mathcal{F}_t)$ -adapted
- $\gamma: \Omega \times [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$  is such that  $(\omega, z) \mapsto \gamma_t(z)$  is  $\mathcal{F}_t \times \mathcal{B}(\mathbb{R})$ -measurable  $\forall t$  and  $t \rightarrow \gamma_t(z)$  is càglàd  $\forall \omega, z$ ;

$$\gamma_t(z)^2 \leq A_t \rho(z), \quad \int_{|z| \leq 1} \rho(z) \nu(dz) < \infty$$

with  $\rho$  positive deterministic and  $A$  càglàd  $(\mathcal{F}_t)$ -adapted.

# Model setup

- The stock price  $S$  is a Lévy-Itô process with coefficients  $\mu, \sigma, \gamma$ ;
- The continuous-time strategy  $\phi$  is a Lévy-Itô process with coefficients  $\tilde{\mu}, \tilde{\sigma}, \tilde{\gamma}$ .
- The agent uses the discrete strategy  $\phi_t^n := \phi_{h\lceil t/h \rceil}$  instead of the continuous strategy  $\phi_t$ .

# The normalizing sequence

The normalizing factor need not be equal to  $\sqrt{n}$ .

Suppose  $\phi$  and  $S$  move only by finite-intensity jumps. If there is only one jump between  $t_i$  and  $t_{i+1}$ ,

$$\int_{t_i}^{t_{i+1}} \phi_{t-} dS_t = \int_{t_i}^{t_{i+1}} \phi_{t-}^n dS_t$$

Therefore  $P[\varepsilon_t^n \neq 0] = O(1/n)$  and

$$n^\alpha \varepsilon_t^n \rightarrow 0$$

in probability  $\forall \alpha$ .

More generally, if  $S$  and  $\phi$  are Lévy-Itô processes without diffusion parts,

$$\sqrt{n} \varepsilon_t^n \rightarrow 0$$

in probability uniformly on  $t$ .

# Main result

Then the discretization error satisfies

$$\begin{aligned}\sqrt{n}\varepsilon_t^n \rightarrow & \sqrt{\frac{T}{2}} \int_0^t \sigma_s \tilde{\sigma}_s dW_s^* + \sqrt{T} \sum_{i: T_i \leq t} \Delta\phi_{T_i} \sqrt{\zeta_i} \xi_i \sigma_{T_i} \\ & + \sqrt{T} \sum_{i: T_i \leq t} \Delta S_{T_i} \sqrt{1 - \zeta_i} \xi_i' \tilde{\sigma}_{T_i-}.\end{aligned}$$

$W^*$  is a standard BM independent from  $W$  and  $J$ ,  
 $(\xi_k)_{k \geq 1}$  and  $(\xi_k')_{k \geq 1}$  are two sequences of independent  $N(0, 1)$ ,  
 $(\zeta_k)_{k \geq 1}$  is sequence of independent  $U([0, 1])$   
 $(T_i)_{i \geq 1}$  are the jump times of  $J$  enumerated in any order.

# Remarks on convergence

- The hedging error  $\sqrt{n}\varepsilon_t^n$  converges weakly in finite-dimensional laws but not in Skorohod topology.
- The discretized error process  $\sqrt{n}\varepsilon_{h[t/h]}^n$  converges in Skorohod topology to the same limit.

# Idea of the proof

Main tool: if  $(X^n)$  and  $(Y^n)$  are two sequences of processes such that

$$\sup_t |X_t^n - Y_t^n| \rightarrow 0 \quad \text{in probability}$$

and  $X^n \rightarrow X$  weakly then  $Y^n \rightarrow X$  weakly.

# Idea of the proof

Step 1 Remove the big jumps

Step 2 Remove the small jumps

Step 3 Now we can write

$$S_t = S_0 + S_t^d + S_t^c + S_t^j$$

$$S_t^d = \int_0^t \left( \mu_s + \int \gamma_s(z) \nu(dz) \right) ds$$

$$S_t^c = \int_0^t \sigma_s dW_s$$

$$S_t^j = \int_0^t \int \gamma_s(z) J(ds \times dz)$$

and  $\phi_t = \phi_0 + \phi_t^d + \phi_t^c + \phi_t^j$ .

# Idea of the proof

The leading terms in the hedging error are

$$\begin{aligned}\sqrt{n} \int (\phi_t^c - \phi_t^{c,n}) dS_t^c &\rightarrow \sqrt{\frac{T}{2}} \int_0^t \sigma_s \tilde{\sigma}_s dW_s^* \\ \sqrt{n} \int (\phi_t^j - \phi_t^{j,n}) dS_t^c &= \sum_i \sqrt{n} \Delta \phi_{T_i} \int_{T_i}^{\psi(T_i)} \sigma_s dW_s \\ &\rightarrow \sqrt{T} \sum_{i: T_i \leq t} \Delta \phi_{T_i} \sqrt{\zeta_i} \xi_i \sigma_{T_i} \\ \sqrt{n} \int (\phi_t^c - \phi_t^{c,n}) dS_t^j &= \sum_i \sqrt{n} \Delta S_{T_i} \int_{\phi(T_i)}^{T_i} \tilde{\sigma}_s dW_s \\ &\rightarrow \sqrt{T} \sum_{i: T_i \leq t} \Delta S_{T_i} \sqrt{1 - \zeta_i} \xi_i' \tilde{\sigma}_{T_i-}.\end{aligned}$$

# Application: delta-hedging in a Lévy market

$$S_t = S_0 e^{X_t}, \quad X_t = bt + \sigma W_t + \int zJ(ds \times dz)$$
$$C(t, S) = E^Q[H(Se^{X_{T-t}})], \quad \phi_t = \frac{\partial C}{\partial S}(t, S_t)$$

Suppose

- The Lévy measure is finite and has a regular density (e.g. Merton model).
- The payoff function  $H$  is piecewise smooth with a finite number of discontinuities.

# Application: delta-hedging in a Lévy market

Apply the Itô formula to get the decomposition for  $\phi$ :

$$d\phi_t = d \frac{\partial C(t, S_t)}{\partial S} = \left\{ \frac{\partial^2 C}{\partial t \partial S} + (b + \sigma^2/2) \frac{\partial^2 C}{\partial S^2} S_t + \frac{\sigma^2}{2} \frac{\partial^3 C}{\partial S^3} S_t^2 \right\} dt + \sigma \frac{\partial^2 C}{\partial S^2} S_t dW_t + \int_{\mathbb{R}} \left( \frac{\partial C}{\partial S}(t, S_{t-} e^z) - \frac{\partial C}{\partial S} C(t, S_{t-}) \right) J(dt \times dz)$$

Under the hypotheses on  $H$  and  $\nu$  it can be shown that the coefficients do not explode in  $T$ : almost all trajectories end in a point where  $H$  is smooth.

# Application: delta-hedging in a Lévy market

The main result then implies  $\sqrt{n}\varepsilon_t^n \rightarrow Z_t$  with

$$\begin{aligned} Z_t = & \sqrt{\frac{T}{2}} \int_0^t \sigma^2 S_s^2 \frac{\partial^2 C}{\partial S^2} dW_s^* + \sqrt{T} \sum \Delta \frac{\partial C}{\partial S} \sqrt{\zeta_i} \xi_i \sigma S_s \\ & + \sqrt{T} \sum \Delta S_s \sqrt{1 - \zeta_i} \xi_i' \sigma S_{s-} \frac{\partial^2 C}{\partial S^2}(s, S_{s-}) \end{aligned}$$

## Application: risk of a hedged option position

If  $E[Z_t^2] < \infty$ , we can estimate the risk of a hedged option position using

$$P[|\epsilon_t^n| \geq \delta] \leq \frac{1}{\delta\sqrt{n}} E[Z_t^2]^{1/2}$$

with (small jump size approximation)

$$E[Z_t^2] \approx \frac{T}{2} \int_0^t E \left[ S_s^4 \left( \frac{\partial^2 C}{\partial S^2} \right)^2 \right] (\sigma^4 + \sigma^2 \int (e^z - 1)^2 (e^{2z} + 1) \nu(dx)).$$

This should be compared to the MSE from market incompleteness:

$$E[\epsilon_t^2] \approx \frac{1}{4} \int_0^t E \left[ S_s^4 \left( \frac{\partial^2 C}{\partial S^2} \right)^2 \right] \int (e^z - 1)^4 \nu(dx).$$

# Application: option hedging with transaction costs

- In presence of transaction costs, the total hedging error comes from the cost of rebalancing (bias) plus the approximation error (variance)
- It makes sense to minimize

$$\frac{1}{n}E[Z_t^2] + E[(C_t^n)^2]$$

where  $C_t^n$  is the transaction cost.

- For proportional transaction cost with  $\Delta C = kS|\Delta\phi|$ ,

$$C_t^n \approx \sqrt{nk} \sqrt{\frac{2}{\pi T}} \int_0^t \sigma S_u^2 \frac{\partial^2 C}{\partial S^2}(u, S_u) du.$$

## Application: option hedging with transaction costs

- Since the transaction cost grows with  $t$  and the hedging error with  $\sqrt{t}$ , the optimal rebalancing step  $h^*$  depends on the length of the holding period.
- For short holding periods,

$$h^* \approx \left( \frac{4tk^2}{\pi(\sigma^2 + \int (e^z - 1)^2)(e^{2z} + 1)\nu(dz)} \right)^{1/2}.$$

- In a Merton model with 10 jumps a year, standard deviation of jump size 5% and a diffusion component volatility of 20%, for a proportional transaction cost of 1% and a holding period of 1 month, this gives  $h^* \approx 4$  days. Without jumps we would have obtained  $h^* \approx 6$  days.

# Extensions

- Other modes of convergence ( $L^2$ ):  
prove that  $E[n\varepsilon_t^2] \rightarrow E[Z_t^2]$ .
- Non-uniform and random rebalancing dates:  
rebalance more often when gamma is big.
- Asymptotic analysis of transaction costs (Leland-Kabanov results) for jump processes.